

Lascar™ Verbena

Uniform series is early to flower. Excellent in combinations.

Bold red color with a distinct dark eye that turns blue in low light conditions.



Lascar™ Vampire Verbena

0	Spread:	Exposure:							
, ,	12 - 16" (30 - 41cm)	Sun							
Rooting Guidelines									
Recommended Cell Size (in millimeters): 25 mm	Hormone (IBA + NAA mix):	Average Days with Mist: 7 to 9	Early PGR Treatmer Recommended (after 10 days): (see note 1) yes	er 7 - Pro	erage ppagation eeks: o 4	Pinch ir Propaga optional		Comments: Avoid stretch; reduce and stop misting as soon as possible.	
Rooting & Finishing Media pH: (see note 2) 5.8-6.2	Fertilizer Needs: (200-250 PPM N/EC 1		ratering: (see note 4) ow on the dry side						
Finishing Guidelines									
Temperature Needs Day/Night: (see note 5)	Pinch after Transplant (unpinched liners)		(PGR) (S) = sp drench:	oray / (D) =	Pests and Fundamental Pests and Pests	Lo , White w	Let media watering.	comments: et media dry out between vatering. Avoid stretch, and	
65-75°F/60-65°F 18- 24°C/16-18°C	yes	5,000 to 6	,000 B-Nine 1,500-2, Sumagic 5-10 P Cycocel 1,000-2	PM (S),	Fly, Leafminer, Powdery Mildew		pinch as needed to improve plant habit.		
Crop Time									

6 to 7 weeks, 1 ppp 7 to 9 weeks, 1 ppp 8 to 10 weeks, 3 to 5 ppp

. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

Start with low concentrations after roots show and increase with plant development to the recommended levels.

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

perature needs day/night

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Lascar™ Varieties



Blue+White Purple+White Purple Black Velvet Orange Lava Burgundy

