

Dahlietta[®] Dahlia

Uniform series for early sales Flowers in as little as 7 weeks Wide range of colors and patterns Best-in-class, easy-to-grow assortment



Dahlietta[®] Anna Dahlia

General & Information	on									
Height:	Spread:	Exposure								
10 - 12" (25 - 30cm)	10 - 12" (25 - 30ci	m) Sun								
Rooting Guidelines										
Recommended Cell	Rooting	Average	Early P	Early PGR Treatment		2	Pinch in		Comments:	
Size (in	Hormone	Days with	ith Recommended (after		Propagation		Propagation:	ion: Hormone will improve		
millimeters):	(IBA + NAA	Mist:	- 10 da	IYS): (see note 1)	Weeks:		optional	Night-break lighting (10 p.m 2 a.m.) will reduce likelihood		
25 mm	mix):	9 to 12	no		3 to 4					
	yes								of tuber formation.	
Rooting & Finishing										
Media pH: (see note 2)	Fertilizer Needs: (see note 3) Watering: (see note 4)									
5.8-6.2	200-250 PPM N/	EC 1.8-2.2	keep eve	keep evenly moist						
Finishing Guidelines										
Temperature	Pinch after	Light L	evels	Plant Growth Reg	ulators	Pests a	and Fungal Diseases:		Comments:	
Needs Day/Night:	Transplant in			(PGR) (S) = spray / (D) =		Phytophthora, Powdery Mildew,		lew,	Night-break lighting will	
(see note 5)	(unpinched lin	ers): Candle	s:	drench:		Botrytis, Pythium, Aphids, Leafminer, Spider Mites, Thrips			reduce likelihood of tuber formation.	
65-76°F/59-64°F 18- 24°C/15-18°C	recommended	4,000 t	6,000	B-Nine 2,500-4,000 PPM (S), Chlormequat chloride 1,000- 1,500 PPM tank mix (S)				rips		

Crop Time

 4- to 5-inch (10- to 13-cm) Pots, Quarts :
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 6 to 8 weeks, 1 ppp
 7 to 9 weeks, 1 to 2 ppp

1. PGR applications during rooting

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Dahlietta[®] Varieties



Patty

Shirley

Candy

Paula

Blanca

Julia



Tessy

Rachel

Emily

Coby

Louise