

Early Love[®] Dianthus

Early flowering, unique bicolor Large, double blooms on short stems Great match to Pink Kisses® Excellent branching Ideal bouquets for Valentine's Day



Early Love® Dianthus

General & Informatio	n									
Height:	Spread:	Exposure:	Hardiness	Zone:						
10 - 14" (25 - 36cm)	8 - 12" (20 - 30cm)	Sun	7a - 7a							
Rooting Guidelines										
Recommended Cell	Rooting	Average	Early PGR Treatment		Average		Pinch in		Comments:	
Size (in	Hormone (IBA +	Days with	Recom	Recommended (after 7 -		Propagation		gation:	Keep very moist for the first 7 days, then reduce	
millimeters):	NAA mix):	Mist:	10 days	S): (see note 1)) Weeks:		recommended			
25/35 mm	recommended	14 to 21	no		5 to 6	0 6			mist to avoid saturated media.	
Rooting & Finishing										
Media pH: (see note 2)	Fertilizer Needs:	(see note 3)	Natering: (see note 4)						
6.2-6.5	100-150 PPM N/EC 1.4-1.8 grow on the dry side									
Finishing Guidelines										
Temperature Needs	Pinch after	Light	Levels	Plant Growth Regu	ators F	Pests and		Comments:		
Day/Night: (see note 5)	Transplant	in Foc	t (PGR) (S) = spray		/(D) Fungal		Remove ea		arly flower buds to	
65-70°F/50-55°F 18-	(unpinched line	rs): Candle	es:	= drench:		Diseases:		promote br	ranching. Cool	
21°C/10-13°C	yes	5,000	0 6,000	Bonzi 1-2 PPM (S)		Spider Mites,		temperatures and high light levels		
·					Fusarium		are ideal.			
Crop Time										

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:10 to 12 weeks (varies seasonally), 1 ppp12 to 14 weeks (varies seasonally), 1 to 2 ppp12 to 14 weeks (varies seasonally), 3 to 5 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.