



## Scully Dianthus

Vivid purple blooms contrast with striking dark foliage  
 Early and mid-Spring sales  
 Versatile for indoor or outdoor use  
 Grow cool for low-energy production  
 Handles cold and wet conditions  
 Thrives in Summer gardens  
 Works well as an indoor potted plant

**selecta** one

Scully Dianthus

| General & Information                      |                                  |  |  |  |                           |   |  |
|--|----------------------------------|--|--|--|---------------------------|---|--|
| Height:                                    | Spread:                          | Exposure:                                      | Hardiness Zone:                                      |  |                           |   |  |
| 10 - 14" (25 - 36cm)                       | 8 - 12" (20 - 30cm)              | Partial Sun, Sun                               | 6a - 6a  |  |                           |   |  |
| Rooting Guidelines                         |                                  |  |  |  |                           |   |  |
| Recommended Cell Size (in millimeters):    | Rooting Hormone (IBA + NAA mix): | Average Days with Mist:                        | Early PGR Treatment Recommended (after 7 - 10 days): | Average Propagation Weeks:                     | Pinch in Propagation:     | Comments:   |  |
| 25/35 mm                                   | recommended                      | 14 to 21                                       | no   | 5 to 6   | recommended               | Keep hydrated. Sensitive root system; avoid overwatering of young plants. |  |
| Rooting & Finishing                        |                                  |  |  |  |                           |   |  |
| Media pH:                                  | Fertilizer Needs:                | Watering:                                      |  |  |                           |   |  |
| (see note 2)                               | (see note 3)                     | (see note 4)                                   |  |  |                           |   |  |
| 6.2-6.5                                    | 100-150 PPM N/EC 1.4-1.8         | grow on the dry side                           |  |  |                           |   |  |
| Finishing Guidelines                       |                                  |  |  |  |                           |   |  |
| Temperature Needs                          | Pinch after                      | Light Levels in                                | Plant Growth Regulators                              | Pests and                                      | Comments:                 |   |  |
| Day/Night:                                 | Transplant                       | Foot Candles:                                  | (PGR) (S) = spray / (D) =                            | Fungal   | Leave early flowers on    |   |  |
| (see note 5)                               | (unpinched liners):              | 5,000 to 6,000                                 | drench:  | Diseases:                                      | plant. Blooms last a long |   |  |
| 55-68°F/45-55°F 13-20°                     | yes                              |  | Bonzi 1-2 PPM (S)                                    | Aphids, Fusarium                               | time.                     |   |  |
| C/7-13°C                                   |                                  |  |  |  |                           |   |  |
| Crop Time                                  |                                  |  |  |  |                           |   |  |
| 4- to 5-inch (10- to 13-cm) Pots, Quarts : |                                  | 6-inch (15-cm) Pots, Gallons:                  |  | 10- to 12-inch (25- to 30-cm) Tubs or Baskets: |                           |   |  |
| 12 to 20 weeks (varies seasonally). 1 ppp  |                                  | 14 to 20 weeks (varies seasonally). 1 to 2 ppp |  |  |                           |   |  |

### 1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

### 2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

### 3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

### 4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

### 5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.