



Dianthus Carmen™ Pink

## Dianthus Carmen™

Zone 5-hardy, true perennial.  
Overwinters better than other carnations.  
Excellent uniformity.  
Double blooms.  
True carnation.  
Blooms continuously until frost.

Double pink flowers bloom continuously – attracting pollinators – until frost.  
Hardy to Zone 5, plants return the following year to bloom again.

**selecta** one

General & Information						
Height:	Spread:	Exposure:	Hardiness Zone:			
10 - 14" (25 - 36cm)	10 - 12" (25 - 30cm)	Partial Sun, Sun	5b - 5b			
Rooting Guidelines						
Recommended Cell Size (in millimeters):	Rooting Hormone (IBA + NAA mix):	Average Days with Mist:	Early PGR Treatment Recommended (after 7 - 10 days):	Average Propagation Weeks:	Pinch in Propagation:	Comments:
25/35 mm	recommended	14 to 21	no	5 to 6	recommended	Keep very moist for the first 7 days, then reduce mist to avoid saturated media.
Rooting & Finishing						
Media pH:	Fertilizer Needs:		Watering:			
6.2-6.5	100-150 PPM N/EC 1.4-1.8		grow on the dry side			
Finishing Guidelines						
Temperature Needs	Pinch after	Light Levels	Plant Growth Regulators	Pests and	Comments:	
Day/Night:	Transplant	in Foot	(PGR) (S) = spray / (D)	Fungal	Remove early flower buds to	
65-70°F/50-55°F 18-21°C/10-13°C	(unpinched liners):	Candles:	= drench:	Diseases:	promote branching. Cool	
	yes	5,000 to 6,000	B-Nine 2,500 PPM (S), Bonzi 2 PPM (S)	Spider Mites, Fusarium	temperatures and high light levels are ideal.	
Crop Time						
4- to 5-inch (10- to 13-cm) Pots, Quarts :		6-inch (15-cm) Pots, Gallons:		10- to 12-inch (25- to 30-cm) Tubs or Baskets:		
12 to 14 weeks (varies seasonally), 1 ppp		14 to 16 weeks (varies seasonally), 1 to 2 ppp		18 to 20 weeks (varies seasonally), 3 to 5 ppp		

### 1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

### 2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

### 3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

### 4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.


### 5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

## Other Carmen™ Varieties



Burgundy

Purple

Red