

DiaDeur[™] Dianthus

Well-branched habit. Low-energy growing; weather tolerant. Fully double, multicolored blooms. Fragrant flowers. Ideal for containers – both indoor and outdoor – and garden use.



DiaDeur™ Magenta+White Edge Dianthus

General & Informati	on					
Height:	Spread:	Exposure:	Hardiness Zone:			
10 - 14" (25 - 36cm)	10 - 12" (25 - 30cm)	Partial Sun, Su	un 6a-6a			
Rooting Guidelines						
Recommended Cell	Rooting	Average	Early PGR Treatment	Average	Pinch in	Comments:
Size (in	Hormone (IBA +	Days with	Recommended (after 7	Propagation	Propagation:	Reduce mist after 4 to 5
millimeters):	NAA mix):	Mist:	- 10 days): (see note 1)	Weeks:	recommended	days to promote root
25/35 mm	recommended	14 to 21	no	5 to 6		formation. Avoid excessive moisture in propagation media.

Rooting & Finishing		
Media pH: (see note 2)	Fertilizer Needs: (see note 3)	Watering: (see note 4)
5.8-6.5	100-150 PPM N/EC 1.4-1.8	grow on the dry side
Finishing Cuidelines		

Finishing Guidelines					
Temperature	Pinch after	Light Levels	Plant Growth Regulators	Pests and	Comments:
Needs Day/Night:	Transplant	in Foot	(PGR) (S) = spray / (D) =	Fungal	Pinch when bud is visible and side
(see note 5)	(unpinched	Candles:	drench:	Diseases:	shoots start to develop. Cool
65-70°F/50-55°F 18-	liners):	5,000 to 6,000	B-Nine 2,500 PPM (S), Cycocel	Spider Mites,	temperatures and high light levels will
21°C/10-13°C	yes		500-750 PPM (S), Bonzi 2-5	Fusarium	improve plant quality.
			PPM are all effective PGRs if		
			needed.		

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts :6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:14 to 20 weeks (varies seasonally), 1 ppp16 to 22 weeks (varies seasonally), 2 ppp16 to 22 weeks (varies seasonally), 4 to 5 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by $8-12^{\circ}$ F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other DiaDeur[™] Varieties



Pink Shades

Magenta+Red

Pink+Red

Velvet+White