



New!

## Sunrise™ Zonal Geranium

Green-leafed, medium-vigor plant  
Large-flowered series blooms early  
Superior color range

**selecta** one

Sunrise™ Light Salmon 26 Zonal Geranium

General & Information						
Height:	Spread:	Exposure:				
12 - 14" (30 - 36cm)	12 - 14" (30 - 36cm)	Sun				
Rooting Guidelines						
Recommended Cell Size (in millimeters): 25/35 mm single or double strip	Rooting Hormone (IBA + NAA mix): optional	Average Days with Mist: 7 to 9	Early PGR Treatment Recommended (after 7 - 10 days): <a href="#">(see note 1)</a> no	Average Propagation Weeks: 4	Pinch in Propagation: no	Comments: Stick right after arrival, keep cuttings hydrated for the first 3 days, and reduce mist afterwards as soon as possible. Apply preventive fungicide treatment after sticking.
Rooting & Finishing						
Media pH: <a href="#">(see note 2)</a> 5.8-6.2	Fertilizer Needs: <a href="#">(see note 3)</a> 200-250 PPM N/EC 1.8-2.0		Watering: <a href="#">(see note 4)</a> grow on the dry side			
Finishing Guidelines						
Temperature Needs Day/Night: <a href="#">(see note 5)</a> 65-75°F/60-65°F 18-24°C/16-18°C	Pinch after Transplant (unpinched liners): no	Light Levels in Foot Candles: 3,000 to 5,000	Plant Growth Regulators (PGR) (S) = spray / (D) = drench: Cycocel 750-1,000 PPM + B-Nine 1,500-2,500 PPM tank mix (S), Sumagic 1-3 PPM (S), Bonzi 2-5 PPM (S)	Pests and Fungal Diseases: Thrips, Fungus Gnats, Spider Mites, White Fly, Rust, Botrytis, Pythium	Comments: Maintain temperature and grow the crop on the dry side, but avoid drying out, as this will slow growth (plants will push flowers). Provide excellent air circulation.	
Crop Time						
4- to 5-inch (10- to 13-cm) Pots, Quarts :		6-inch (15-cm) Pots, Gallons:		10- to 12-inch (25- to 30-cm) Tubs or Baskets:		
8 to 12 weeks, 1 ppp		12 to 14 weeks, 1 ppp		12 to 14 weeks, 3 to 4 ppp		

### 1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

### 2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

### 3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

### 4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.


### 5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.


The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.


## Other Sunrise™ Varieties




Bright Scarlet 26




Coral




Pink




Appleblossom



Ruby + Fringe




White + Zest



Purple + Big Eye




Lavender




Rose + Big Eye




Red




Hot Pink




Bright Lilac




Brilliant Red




Hot Rose + Eye



Orange




Salmon




Violet



White



Dark Red



Light Pink