

# Sunrise<sup>™</sup> Zonal Geranium

Green-leafed, medium-vigor plant Large-flowered series blooms early Superior color range

This new big-eye variety has pink and red bicolor blooms.



Sunrise™ Purple+Big Eye Zonal Geranium

| General & Informati                           | ion                              |                               |  |                                       |                                |  |
|---|----------------------------------|-------------------------------|--|---------------------------------------|--------------------------------|--|
| Height:<br>12 - 14" (30 - 36cm)               | Spread:<br>12 - 14" (30 - 36     | Exposu<br>Scm) Sun            | re:  |                                       |                                |  |
| Rooting Guidelines                            |                                  |                               |  |                                       |                                |  |
| Recommended<br>Cell Size (in<br>millimeters): | Rooting<br>Hormone<br>(IBA + NAA | Average<br>Days with<br>Mist: | Early PGR Treatment<br>Recommended (after<br>7 - 10 days): (see note | Average<br>Propagation<br>Weeks:<br>4 | Pinch in<br>Propagation:<br>no | Comments:<br>Stick right after arrival, keep<br>cuttings hydrated for the first 3<br>days, and reduce mist afterwards<br>as soon as possible. Apply<br>preventive fungicide treatment<br>after sticking. |
| 25/35 mm single or<br>double strip            | mix):<br>optional                | 7 to 9                        | 1)<br>no   |                                       |                                |  |

#### Rooting & Finishing Media pH: (see note 2) Fertilizer Needs: ( Watering: (see note 4) 5.8-6.2 200-250 PPM N/EC 1.8-2.0 grow on the dry side Finishing Guidelines Light Levels Pinch after Plant Growth Regulators Pests and Fungal Temperature Comments: (PGR) (S) = spray / (D) = Needs Day/Night: Maintain temperature and grow the Transplant in Foot Diseases: Candles crop on the dry side, but avoid drying (unpinched drench: Thrips, Fungus Cycocel 750-1,000 PPM + B-Gnats, Spider Mites, 65-75°F/60-65°F 18-3.000 to out, as this will slow growth (plants will liners): Nine 1,500-2,500 PPM tank push flowers). Provide excellent air 5 000 White Fly, Rust, 24°C/16-18°C no mix (S), Sumagic 1-3 PPM (S), Bonzi 2-5 PPM (S) Botrytis, Pythium circulation.

Crop Time

 4- to 5-inch (10- to 13-cm) Pots, Quarts:
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 8 to 12 weeks, 1 ppp
 12 to 14 weeks, 1 ppp
 12 to 14 weeks, 3 to 4 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

### 2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

#### 4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

## Other Sunrise<sup>™</sup> Varieties



1

Rose+Big Eye



Hot Pink

Bright Lilac



White+Zest

Lavender



Orange

Salmon

Violet

White



Dark Red





Hot Rose+Eye