

Super Moon[™] Zonal Geranium

Large-flowered zonal Vibrant blooms Very dark stems provide contrast

Large-flowered zonal geranium features vibrant red blooms on a strong, vigorous plant. Dark stems provide contrast in containers. Heat tolerance makes it ideal for landscapes.



Super Moon™ Red Zonal Geranium

General & Information	n							
Height:	Spread:	Exposure:						
12 - 18" (30 - 46cm)	10 - 14" (25 - 36cm)	Partial Sun, Sun						
Rooting Guidelines								
Recommended Cell	Rooting	Average	e Early PGR Treatment		Ave	rage	Pinch in	Comments:
Size (in millimeters):	Hormone (IBA +	Days with Recon		mended (after 7 - 10	after 7 - 10 Prop		Propagation:	Cycocel 750
30 mm	NAA mix):	Mist:	days): (see note 1)		Wee	ks:	no	PPM spray.
	yes	7 to 9	yes		5			
Rooting & Finishing								
Media pH: (see note 2)	Fertilizer Needs: (see note 3) Watering: (see note 4)							
5.8-6.2	200-250 PPM N/EC 1.8-2.0 grow on the dry side							
Finishing Guidelines								
Temperature Needs	Pinch after	Light Le	ight Levels in Plant Growth Regulat		ors	ors Pests and Fungal Disease		Comments:
Day/Night: (see note 5)	Transplant	plant Foot Car		dles: (PGR) (S) = spray / (I		D) = Botrytis, Pythium, Ralstonia,		
65-77°F/55-65°F 18-25°			7,000	drench:		Xanthomonas, Mites,		
C/13-18°C			Cycocel 700-1,500 PPM	(S) Worms/Caterpillars		pillars		
Crop Timo								

Crop Time

4- to 5-inch (10- to 13-cm) Pots, Quarts:6-inch (15-cm) Pots, Gallons:10- to 12-inch (25- to 30-cm) Tubs or Baskets:7 to 10 weeks, 1 ppp8 to 12 weeks, 1 ppp10 to 14 weeks, 1 to 3 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

5. Temperature needs day/night

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Super Moon[™] Varieties



Bright Red

Violet