

## Magadi™ Compact Lobelia

High Density.

Early-flowering series features brilliant colors Densely branched



Magadi™ Compact Dark Blue Lobelia

Rooting Guidelines         Recommended Cell       Rooting       Average       Early PGR Treatment       Average       Pinch in       Comments:         Millimeters):       + NAA mix):       Mist:       10 days): (see note 1)       Weeks:       Propagation       Keep cuttings         25 mm       no       7 to 9       yes       4       Propagation:       Keep cuttings         Media pH: (see note 2)       Fertilizer Needs:       (see note 3)       Watering: (see note 4)       grow on the dry side         Finishing Guidelines       Pinch after       Light Levels       Plant Growth Regulators       Pests and Fungal       Comments:         70-75°F/55-60°F 21-       24°C/13-16°C       9es       5,000 to 6,000       B-Nine 1,500-2,500 PPM (D)       Pests and Fungal       Comments:         24°C/13-16°C       yes       5,000 to 6,000       B-Nine 1,500-2,500 PPM (D)       Botypis, Pythium       Spider Mites, Botypis, Pythium	5	n pread: - 12" (20 - 30cm)	Exposure: Partial Sun							
Rooting & Finishing         Media pH: (see note 2)       Fertilizer Needs: (see note 3)       Watering: (see note 4)         5.8-6.2       200-250 PPM N/EC 1.8-2.0       grow on the dry side         Finishing Guidelines       Pinch after       Light Levels       Plant Growth Regulators       Pests and Fungal       Comments:         Day/Night: (see note 5)       Pinch after       Light Levels       Plant Growth Regulators       Diseases:       Pinch as needed for improved         70-75°F/55-60°F 21-       (unpinched liners):       Candles:       drench:       Thips, Aphids,       plant habit. Let media dry out between watering.         24°C/13-16°C       yes       5,000 to 6,000       B-Nine 1,500-2,500 PPM (S), Sumagic 3-5 PPM       Botrytis, Pythium	Recommended Cell Size (in millimeters):	Hormone (IBA + NAA mix):	Days with Mist:	n Recon 10 day	nmended (after 7 -	Propag Weeks	gation	Propagati		Keep cuttings hydrated. Avoid overwatering of young
Temperature Needs Day/Night: (see note 5)Pinch after TransplantLight Levels in FootPlant Growth Regulators (PGR) (S) = spray / (D) = drench:Pests and Fungal Diseases:Comments: Pinch as needed for improved plant habit. Let media dry out between watering.70-75°F/55-60°F 21- 24°C/13-16°C(unpinched liners): vesCandles: 5,000 to 6,000drench: B-Nire 1,500-2,500 PPM (S), Sumagic 3-5 PPMPests and Fungal Diseases:Comments: Pinch as needed for improved plant habit. Let media dry out between watering.	Media pH: (see note 2)									·
	Temperature Needs Day/Night: (see note 5) 70-75°F/55-60°F 21-	Transplant (unpinched liners):		oot ndles:	(PGR) (S) = spray drench: B-Nine 1,500-2,500 Sumagic 3-5 PPM	/ (D) = PPM (S),	Diseases Thrips, Ap Spider Mite	hids, plant l es, betwe		as needed for improved nabit. Let media dry out

 4 - to 5-inch (10- to 13-cm) Pots, Quarts :
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 6 to 8 weeks, 1 ppp
 8 to 10 weeks, 1 to 2 ppp
 8 to 10 weeks, 4 to 5 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

## 3. Fertilizer needs

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

## Other Magadi<sup>™</sup> Compact Varieties



Blue Bay

White

Purple

Blue