

## Headliner™ Petunia

Early flowering for a fast finish Dramatic color range on large blooms Wide assortment of Sky petunias

Long-lasting red flowers bloom early for a fast finish. Versatile, nicely mounded Headliner petunias can be planted in gardens, containers and baskets.



Headliner™ Crimson Petunia

Recommended Rooting Average Early PGR Average Pinch in Propagation: Avoid stretch; apply preventive fungicide spray after sticking. For detailed culture on the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petunias, visit SelectaNorthAmerica.com/SkyPetuniaCulture sembles and propagation of the Sky Petuni	Height: 10 - 16" (25 - 41cm)	Spread: 20 - 30" (5		xposure: Gun				
ell Size (in Hormone Days Treatment Recommended (IBA + with NAA mix): Mist: (after 7 - 10 days): (see note 1) yes  Rooting & Finishing Hedia pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4)	Rooting Guidelines							
Media pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4)  4-5.8 250-300 PPM N/EC 2.0-2.4 keep constantly moist  inishing Guidelines  emperature Needs emperature Needs Pinch after Light Levels In Foot (PGR) (S) = spray / (D) = Diseases: Cool temperatures and high (unpinched liners): Candles: drench: Aphids, Thrips, light levels will improve plant (unpinched liners): Pests and Fungal Comments: Cool temperatures and high (unpinched liners): Candles: drench: Aphids, Thrips, light levels will improve plant (S)-75°F/50-60°F 18-4°C/10-16°C Sumagic 20-30 PPM (S), Powdery Mildew, (S)/Sumagic 0.5 PPM (D), Pythium	Cell Size (in Hamillimeters): (25 mm	Hormone Days (IBA + with NAA mix): Mist:		Treatment Recommended (after 7 - 10 days): (see note 1)	Propagation Weeks: no 4		Avoid stretch; apply preventive fungicide spray after sticking. For detailed culture on the Sky Petunias, visit	
emperature Needs Pinch after Transplant in Foot (PGR) (S) = spray / (D) = Diseases: Cool temperatures and high light levels will improve plant for tench: 5-75°F/50-60°F 18- 4°C/10-16°C   Fig.   Comments: Cool temperatures and high light levels will improve plant for the provided from the provided fr	Media pH: (see note 2) 5.4-5.8	250-300 I	,	,				
	Femperature Needs Day/Night: (see note 5) 55-75°F/50-60°F 18- 24°C/10-16°C	Pinch af Transpla (unpinch	nnt ned liners):	in Foot Candles:	(PGR) (S) = spray drench: B-Nine 2,500 PPM (S Sumagic 20-30 PPM (S)/Sumagic 0.5 PPM	/ (D) = D A <sub>I</sub> ), Le	iseases: phids, Thrips, eafminer, White Fly, pwdery Mildew,	Cool temperatures and high light levels will improve plant

6 to 7 weeks, 1 ppp 7 to 9 weeks, 1 ppp 9 to 12 weeks, 3 to 4 ppp

# 1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

Best results are achieved if media with the correct pH is used starting with propagation.

Start with low concentrations after roots show and increase with plant development to the recommended levels.

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

### Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

### Other Headliner™ Varieties















Sky Blue