

Lascar[™] Verbena

Uniform series is early to flower Excellent in combinations



Lascar™ Black Velvet Verbena

n								
Spread:	Exposure:							
12 - 16" (30 - 41cm)	Sun							
Rooting	Average	Early PGR Treatment	Aver	age	Pinch in		Comments:	
· ·	Days with	Recommended (after 7 -		eks: optional		gation:	Avoid stretch; reduce and stop misting as	
+ NAA mix):	Mist:	10 days): (see note 1)						
optional	7 to 9	yes	3 to 4				soon as possible.	
Fertilizer Needs:	(see note 3) W	atering: (see note 4)						
200-250 PPM N/EC	1.8-2.0 gr	ow on the dry side						
emperature Needs Pinch after Day/Night: (see note Transplant		els Plant Growth Regula	ators	Pests and Fu	0		nts:	
		(PGR) (S) = spray /	(D) =	Diseases:			lia dry out between	
(unpinched liners	s): Candles:	drench:		Thrips, Aphids,	White waterin		g. Avoid stretch, and	
yes	5,000 to 6	Sumagic 5-10 PPM (S)		Fly, Leafminer, Powdery Mildew		pinch as needed to improve plant habit.		
	12 - 16" (30 - 41cm) Rooting Hormone (IBA + NAA mix): optional Fertilizer Needs: 200-250 PPM N/EC Pinch after Transplant (unpinched liners	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Average Hormone (I BA Days with + NAA mix): Mist: optional 7 to 9 Fertilizer Needs: (see note 3) W 200-250 PPM N/EC 1.8-2.0 gr Pinch after Light Lev Transplant in Foot (unpinched liners): Candles:	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Hormone (I BA + NAA mix): Average Mist: Early PGR Treatment Recommended (after 7 - 10 days): (see note 1) optional 7 to 9 yes Fertilizer Needs: (see note 3) Watering: (see note 4) 200-250 PPM N/EC 1.8-2.0 grow on the dry side Pinch after Transplant Light Levels in Foot Plant Growth Regula (PGR) (S) = spray / drench: yes 5,000 to 6,000 B-Nine 1,500-2,500 PF Sumagic 5-10 PPM (S)	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Hormone (IBA optional Average Days with Nist: Early PGR Treatment Recommended (after 7 - 10 days): (see note 1) Aver Prop Week 3 to 4 Fertilizer Needs: (see note 3) Watering: (see note 4) 3 to 4 200-250 PPM N/EC 1.8-2.0 grow on the dry side Plant Growth Regulators (PGR) (S) = spray / (D) = (unpinched liners): Candles: yes 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S),	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Average Hormone (IBA Days with Propagation Weeks: optional 7 to 9 Yerage yes String 3 to 4 Pertilizer Needs: (see note 1) 200-250 PPM N/EC 1.8-2.0 Watering: (see note 4) grow on the dry side grow on the dry side Pinch after Light Levels Plant Growth Regulators (unpinched liners): Candles: drench: yes 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S), Sumagic 5-10 PPM (S), Fiy, Leafminer, Powdery Mildee	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Hormone (IBA potional Average Days with Mist: Early PGR Treatment Recommended (after 7 - yes Average Propagation yes Pinch ir Propagation Fertilizer Needs: (see note 3) 200-250 PPM N/EC 1.8-2.0 Watering: (see note 4) grow on the dry side 3 to 4 Pinch after Transplant Light Levels in Foot Plant Growth Regulators (PGR) (S) = spray / (D) = (unpinched liners): Pests and Fungal Diseases: Thrips, Aphids, White Yes yes 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S), Sumagic 5-10 PPM (S), Pests and Fungal Diseases: Thrips, Aphids, White	Spread: Exposure: 12 - 16" (30 - 41cm) Sun Rooting Hormone (IBA potional Average Days with Nist: Early PGR Treatment Recommended (after 7 - yes Average Propagation 3 to 4 Pinch in Propagation: optional Fertilizer Needs: (see note 3) Watering: (see note 4) grow on the dry side 3 to 4 Pinch in Propagation: Pinch in Propagation: Pinch after Transplant Light Levels in Foot Plant Growth Regulators (PGR) (S) = spray / (D) = (unpinched liners): Pests and Fungal Diseases: Thrips, Aphids, White yes Comment Let medi watering pinch as habit.	

Crop Time

 4- to 5-inch (10- to 13-cm) Pots, Quarts:
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 6 to 7 weeks, 1 ppp
 7 to 9 weeks, 1 ppp
 8 to 10 weeks, 3 to 5 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Lascar[™] Varieties



Red+Eye

Blue+White

Orange Lava

Purple+White

Vampire

Burgundy



Dark Red

Dark Violet

Mango Orange

Pink

White

Red Rose