

Blues[™] Verbena

Big blooms with an eye Perfect in mixed containers High resistance to powdery mildew



Blues™ Violet+Eye Verbena

| n | | | | | | | |
|--|--|---|---|--|--|---|--|
| Spread: | Exposure: | | | | | | |
| 12 - 18" (30 - 46cm) | Sun | | | | | | |
| | | | | | | | |
| Rooting Hormone (IBA + NAA mix): optional | Average Days with Mist: 7 to 9 | Early PGR Treatment Recommended (after 7 - 10 days): (see note 1) ves | Prop Wee | pagation eks: | Pinch in Propagation: optional | | Comments: Avoid stretch; reduce and stop misting as soon as possible. |
| | | 0 | | | | | |
| | | | | | | | |
| Pinch after Transplant (uppinched | in Foot | (PGR) (S) = spray / | | Pests and Fungal Diseases: Thrips, Aphids, White Fly, Leafminer, Powdery Mildew | | Comments: Let media dry out between watering. Avoid stretch, and pinch as needed to improve plant habit. | |
| liners): yes | | 000 B-Nine 1,500-2,500 PF | l, | | | | |
| | Spread: 12 - 18" (30 - 46cm) Rooting Hormone (IBA + NAA mix): optional Fertilizer Needs: 200-250 PPM N/EC Plinch after Transplant (unpinched liners): | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Average Hormone (IBA Days with + NAA mix): Mist: optional 7 to 9 Fertilizer Needs: (see note 3) W 200-250 PPM N/EC 1.8-2.0 gr Pinch after Light Lev Transplant in Foot (unpinched Candles: 5,000 to 6, | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Average Early PGR Treatment Hormone (IBA Days with Recommended (after 7 - + NAA mix): Mist: 10 days): (see note 1) optional 7 to 9 yes Fertilizer Needs: (see note 3) Watering: (see note 4) 200-250 PPM N/EC 1.8-2.0 grow on the dry side Pinch after Light Levels Plant Growth Regular Transplant in Foot (PGR) (S) = spray / (unpinched Candles: drench: liners): 5,000 to 6,000 B-Nine 1,500-2,500 PP yes Sumagic 5-10 PPM (S) | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Average Early PGR Treatment Average Hormone (IBA Days with Recommended (after 7 - Programment) Wee optional 7 to 9 yes 3 to Fertilizer Needs: (see note 3) Watering: (see note 4) 3 to 200-250 PPM N/EC 1.8-2.0 grow on the dry side 9 Pinch after Light Levels Plant Growth Regulators Transplant in Foot (PGR) (S) = spray / (D) = (unpinched Liners): 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S), | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Average Early PGR Treatment Average Hormone (IBA Days with Recommended (after 7 - Propagation + NAA mix): Mist: 10 days): (see note 1) Weeks: optional 7 to 9 yes 3 to 4 Fertilizer Needs: (see note 3) Watering: (see note 4) 3 to 4 200-250 PPM N/EC 1.8-2.0 grow on the dry side Plant Growth Regulators Pests and Fu Pinch after Light Levels Plant Growth Regulators Pests and Fu Transplant in Foot (PGR) (S) = spray / (D) = Diseases: (unpinched Candles: drench: Thrips, Aphids Jiners): 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S), Fly, Leafminer yes Sumagic 5-10 PPM (S), Powdery Milde | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Hormone (IBA + NAA mix): Average Mist: Early PGR Treatment Recommended (after 7 - yes Average Propagation Pinch in Propagation optional 7 to 9 yes 3 to 4 Fertilizer Needs: (see note 3) Watering: (see note 4) 3 to 4 200-250 PPM N/EC 1.8-2.0 grow on the dry side Plant Growth Regulators (unpinched liners): Pests and Fungal Diseases: Thrips, Aphids, White | Spread: Exposure: 12 - 18" (30 - 46cm) Sun Rooting Hormone (IBA + NAA mix): Average Mist: Early PGR Treatment Recommended (after 7 - yes Average Propagation Pinch in Propagation: optional 7 to 9 yes 3 to 4 Fertilizer Needs: (see note 3) Watering: (see note 4) optional 200-250 PPM N/EC 1.8-2.0 grow on the dry side Plant Growth Regulators in Foot Pests and Fungal (PGR) (S) = spray / (D) = Pests and Fungal Diseases: Comme Light Levels Pinch after Transplant (unpinched Iners): Light Levels 5,000 to 6,000 B-Nine 1,500-2,500 PPM (S), 5,000 to 6,000 Pests and Fungal Thrips, Aphids, White Comme pinch as |

 4- to 5-inch (10- to 13-cm) Pots, Quarts :
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 6 to 7 weeks, 1 ppp
 7 to 9 weeks, 1 ppp
 8 to 10 weeks, 3 to 4 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Blues[™] Varieties



White

Red

Lavender+Eye

Light Pink+Eye

Magenta+Eye

Purple+Eye



Rose+Eye