

Beats™ Verbena

Flowers 2 weeks earlier than other verbena.

Tolerant to heat, cycling and water stress.

No PGRs; no pinch.

Pink blooms with a white eye cover this naturally compact, spreading and earlyflowering verbena from Spring through Summer.



Beats™ Magenta Verbena

General & Informa	tion					
Height: 6 - 8" (15 - 20cm)	Spread: 10 - 12" (25 - 30cm)	Exposure: Sun				
Rooting Guidelines						
Recommended Cell Size (in millimeters): 25 mm	I Rooting Hormone (IBA + NAA mix): optional	Average Days with Mist: 7 to 9	Early PGR Treatment Recommended (after 7 10 days): (see note 1) no	Average 7 - Propagat Weeks: 2 to 3	Pinch in ion Propagati optional	on: Comments: Avoid overwatering after root formation.
Rooting & Finishing Media pH: (see note 2 5.8-6.2			atering: (see note 4) ow on the dry side			
Finishing Guideline	S					
Temperature Need		Light Lev		Plant Growth Regulators (PGR)		Comments:
Day/Night: (see note 5)	 Transplant (unpinched liner 	in Foot s): Candles:	(S) = spray / (D) Very little or no PGF	R is necessary for	Fungal Diseases: Thrips, Aphids,	Due to its natural compact habit, use care to avoid
65-75°F/60-65°F 18- 24°C/16-18°C	optional	5,000 to 6	,000 Beats. If necessary, PPM/Cycocel 500 PF		Leafminer, Powdery Mildew	overwatering.
Crop Time						
4- to 5-inch (10- to	13-cm) Pots, Quarts	: 6-inch (15-	cm) Pots. Gallons: 10-	to 12-inch (25- to	o 30-cm) Tubs or Ba	skets:

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

7 to 9 weeks, 5 ppp

. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs

5 to 6 weeks, 1 ppp

Start with low concentrations after roots show and increase with plant development to the recommended levels.

6 to 8 weeks, 3 ppp

Watering

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein

Other Beats[™] Varieties



Blue+White Bright Purple Blue Purple+White Pink Shades Red





Red+White

White