

MiniFamous[®] Neo Calibrachoa

Uniform habit Large, single and double blooms Exclusive double bicolors



MiniFamous[®] Neo White Calibrachoa

General & Informat	ion						
Height:	Spread:	Exposure:					
8 - 12" (20 - 30cm)	12 - 15" (30 - 38cm)	Sun					
Rooting Guidelines							
Recommended Cell	Rooting	Average	Early PGR Treatment	Average	Pinch in	Comments:	
Size (in	Hormone (IBA	Days with	Recommended (after 7 -	Propagation	Propagation:	Don't stick too deep;	
millimeters):	+ NAA mix):	Mist:	10 days): (see note 1)	Weeks:	optional	avoid overwatering of	
25 mm	optional	9 to 12	yes	4 to 5		young plants and water stress.	
Rooting & Finishing							
Media pH: (see note 2	edia pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4)						
5.4-5.8	200-250 PPM N/EC	2.0-2.4 ke	ep evenly moist				
Finishing Guidelines	5						
Temperature	Pinch after	Light Levels	Plant Growth Regulators	Pests and Funga	al Commer	its:	
Needs Day/Night:	Transplant	in Foot	(PGR) (S) = spray / (D)	Diseases:	Start Cali	brachoa warm. After side	
(see note 5)	(unpinched	Candles:	= drench:	Powdery Mildew,	shoots ha	ve developed, cool	
70-75°F/55-60°F 21-	liners):	5,000 to	B-Nine 2,500-3,500 PPM	Pythium, Thielavic	psis, temperat	temperatures and high light levels will	
24°C/13-16°C	recommended 6,000	(S), Sumagic 10-20 PPM (S)/Sumagic 0.25 PPM (D), Bonzi 1-2 PPM (D)	TMV, Aphids, Fung Gnats, Thrips		improve plant quality. Watch for Fe deficiency.		

 Crop Time

 4- to 5-inch (10- to 13-cm) Pots, Quarts :
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 7 to 8 weeks, 1 ppp
 7 to 11 weeks, 1 to 2 ppp
 9 to 12 weeks, 4 to 5 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots

have developed nicely. Afterward follow recommendations. The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges).

F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other MiniFamous[®] Neo Varieties



Magenta Star

Rose Flash

Double Red

Double PlumTastic

Red Shuffle

Dark Blue

Light Pink+Eye



Deep Orange











Coral Flash Red Hawaii



Double Dark Blue

Violet I ce



Double Deep Yellow



Double OrangeTastic









Double Amethyst



Double Purple



Vampire

Orange+Red Eye

Pink

Salmon Pink+Eye

Blue