

MiniFamous® Neo Calibrachoa

Uniform habit Large, single and double blooms Exclusive double bicolors

Bicolor eye pattern on a double flower – a pattern unique to Selecta One! Same color as Uno PlumTastic on the Neo habit, for a more vigorous, trailing variety. Matches Neo OrangeTastic in vigor and timing.



MiniFamous® Neo Double PlumTastic

Recommended Cell Rooting Average Hormone (IBA Days with Recommended (after 7 - Propagation Propagation: Don't stick too deep; avoid overwatering of young plants and water stress. Rooting & Finishing Media pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4) keep evenly moist Finishing Guidelines Temperature Pinch after Needs Day/Night: Transplant (unpinched Candles: = drench: (unpinched Candles: = drench: powdery Mildew, pythium, Thielavlopsis, temperatures and high light levels water and representations. Rooting & Finishing Watering: (see note 4) keep evenly moist Finishing Guidelines Fertilizer Needs: (see note 3) Watering: (see note 4) keep evenly moist Finishing Guidelines For young plants and water stress. Pests and Fungal Comments: Start Calibrachoa warm. After side showed warm.		on Spread: 12 - 15" (30 - 38cm)	Exposure: Sun				
Size (in Hormone (IBA Days with NMst: 10 days): (see note 1) Weeks: optional avoid overwatering of young plants and water stress. Rooting & Finishing Media pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4) keep evenly moist Finishing Guidelines Temperature Needs Day/Night: Transplant in Foot (PGR) (S) = spray / (D) (see note 5) (unpinched (Lipht Levels Candles: = drench: provided of the content of t	Rooting Guidelines						
Media pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4) keep evenly moist Finishing Guidelines Femperature Pinch after Light Levels in Foot (PGR) (S) = spray / (D) Diseases: Start Calibrachoa warm. After side (unpinched (unpinched Candles: = drench: Powdery Mildew, procypitals): Tecommended 6,000 (S), Sumagic 10-20 PPM TMV, Aphids, Fungus Improve plant quality. Watch for Fe (S)/Sumagic 0.25 PPM (D), Gnats, Thrips deficiency.	Recommended Cell Size (in millimeters): 25 mm	Hormone (IBA + NAA mix):	Days with Mist:	Recommended (after 7 - 10 days): (see note 1)	Propagation Weeks:	Propagation:	Don't stick too deep; avoid overwatering of young plants and water
Temperature Pinch after Light Levels in Foot (PGR) (S) = spray / (D) Diseases: Start Calibrachoa warm. After side (see note 5) (unpinched (207-75°F/55-60°F 21- liners): 5,000 to B-Nine 2,500-3,500 PPM (D), Candles: ecommended (S)/Sumagic 0.25 PPM (D), Gnats, Thrips (deficiency.	Rooting & Finishing Media pH: (see note 2) 5.4-5.8	200-250 PPM N/EC					
	Finishing Guidelines Temperature Needs Day/Night: (see note 5) 70-75°F/55-60°F 21- 24°C/13-16°C	Pinch after Transplant (unpinched liners):	in Foot Candles: 5,000 to	(PGR) (S) = spray / (D) = drench: B-Nine 2,500-3,500 PPM (S), Sumagic 10-20 PPM (S)/Sumagic 0.25 PPM (D),	Diseases: Powdery Mildew, Pythium, Thielavio TMV, Aphids, Fung	Start C shoots psis, temper jus improv	alibrachoa warm. After side have developed, cool atures and high light levels wi e plant quality. Watch for Fe

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

Fertilizer needs:

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering:

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

5. Temperature needs day/night:

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by 8-12° F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other MiniFamous® Neo Varieties



Double Amethyst