

Bounce[™] Interspecific Impatiens

Best for shade and heat Loads of color in the garden Great branching; fast flowering Downy mildew resistant



Bounce[™] White Interspecific Impatiens

General & Information	on								
Height:	Spread:	Exposure:							
14 - 20" (36 - 51cm)	14 - 20" (36 - 51cm)	Partial Sun							
Rooting Guidelines									
Recommended Cell	Rooting	Average	Early PGR Treatment	Av	Average	Pinch in	h in		
Size (in	Hormone (IBA	Days with	Recommended (after 7 -		opagation	Prop	pagation:		
millimeters):	+ NAA mix):	Mist:	10 days): (see note 1)		eeks:	no		starting day 5; don't overwater.	
25 mm	no	6 to 9	yes	4 t	:0 5				
Rooting & Finishing									
Media pH: (see note 2) Fertilizer Needs: (see note 3) Watering: (see note 4)									
6.2-6.5	100-150 PPM N/EC 1.4-1.8 grow on the dry side								
Finishing Guidelines									
Temperature Needs	Pinch after	Light Lev	els Plant Growth		Pests and Fungal Diseases: Thrips, Aphids, Mites, INSV, Botrytis, Pythium		Comments		
Day/Night: (see note	Transplant	in Foot	Regulators (PGR) (Don't overwater or overfeed.		
5)	(unpinched liner	,	spray / (D) = drer					ove 6 to avoid	
68-79°F/55-64°F 20- 26°C/13-18°C	no	4,000 to 9	,000 B-Nine 1,500 PPM (S)			iron/manganese toxicity.		
Crop Time									

 4- to 5-inch (10- to 13-cm) Pots, Quarts :
 6-inch (15-cm) Pots, Gallons:
 10- to 12-inch (25- to 30-cm) Tubs or Baskets:

 6 to 8 weeks, 1 ppp
 8 to 10 weeks, 1 ppp
 10 to 13 weeks, 3 to 4 ppp

1. PGR applications during rooting:

Sprays only. Cover the plants and avoid run-off as much as possible. For vigorous products, first application should be done about 7-10 days after sticking, second when needed. Please test before use.

2. Media pH:

Best results are achieved if media with the correct pH is used starting with propagation.

3. Fertilizer needs

Start with low concentrations after roots show and increase with plant development to the recommended levels.

4. Watering

In general avoid drying out or overwatering the substrate as this will affect plant quality and finishing time. Root system will suffer when conditions change from one extreme to the other. "Growing on the dry side" means letting the substrate become dry (substrate color turns light brown and substrate starts to separate from the pot edge) before watering.

"Keep constantly moist" means watering the plants just before substrate becomes dry. Water management in rooting and finishing is half the success and the best way to prevent root-bound diseases, improve performance and control plant quality.

During rooting keep soil temperature measured in the liner at 70°F/21°C until cuttings are well-rooted. After planting the liner, keep temperature up until roots have developed nicely. Afterward follow recommendations.

The use of "Cool Morning" treatments will improve plant quality, flower and foliage color, and plant habit, and harden the plants. Lower the temperature by $8-12^{\circ}$ F/4-6°C 2 hours before sunrise until 2-3 hours after sunrise. Start treatments after plants are rooted in, well-branched and cover the media (reach the pot edges). To avoid delayed flowering, keep the average day temperature under a close watch and if necessary, increase night temperatures. Finishing of the crop depends on the average day temperature, and positive or negative DIF methods as well as "Cool Morning" can be used to improve plant quality.

NOTE: Growers should use the information presented here as guidelines only. Selecta recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by Selecta of any products listed herein. Selecta's terms and conditions of sale shall apply to all products listed herein.

Other Bounce[™] Varieties



Bright Coral

Violet

Cherry

Pink Flame